Molecular Brain Research, 9 (1991) 359–360
Elsevier

Author Index

Augood, S.J., Kiyama, H., Faull, R.L.M. and Emson, P.C., Differential effects of acute dopaminergic D_1 and D_2 receptor antagonists on proneurotensin mRNA expression in rat striatum, 341

Barden, N., see l'Héreault, S., 327
Barg, J., see Coscia, C.J., 299
Barnard, E.A., see Bateson, A.N., 333
Barnard, E.A., see Coscia, C.J., 299
Bartlett, W.P., see Li, X.-s., 313
Barton, A.J.L., see Harrison, P.J., 15
Barton, A.J.L., see Harrison, P.J., 161
Bateson, A.N., Harvey, R.J., Wisden, W., Glencorse, T.A., Hicks, A.A., Hunt, S.P., Barnard, E.A. and Darlison, M.G., The chicken $GABA_A$ receptor $\alpha 1$ subunit: cDNA sequence and localization of the corresponding mRNA, 333
Belcheva, M.M., see Coscia, C.J., 299
Bem, W.T., see Coscia, C.J., 299
Benson, D.L., Isackson, P.J. and Jones, E.G., In situ hybridization reveals VIP precursor mRNA-containing neurons in monkey and rat neocortex, 169
Bergeron, C., see Sommerville, M.J., 1
Bertolucci, P.H.F., see Harrison, P.J., 161
Beyreuther, K., see König, G., 259
Bierer, L., see Parks, K.M., 125
Bowen, D.M., see Harrison, P.J., 161
Bramson, H.N., see Slemmon, J.R., 245
Brodin, E., see Srinivasan, M., 233

Calignano, A., Piomelli, D., Sacktor, T.C. and Scheartz, J.H., A phospholipase A_2-stimulating protein regulated by protein kinase C in *Aplysia* neurons, 347
Campbell, G.A., see Slemmon, J.R., 245
Carver, C.H., see Kim, K.S., 277
Caviccholi, L., Flanigan, T.P., Dickson, J.G., Vantini, G., Toso, R.D., Fusco, M., Walsh, F.S. and Leon, A., Choline acetyltransferase messenger RNA expression in developing and adult rat brain: regulation by nerve growth factor, 319
Ceccatelli, S., Cortés, R. and Hökfelt, T., Effect of reserpine and colchicine on neuropeptide mRNA levels in the rat hypothalamic paraventricular nucleus, 57
Chao, H.M. and McEwen, B.S., Glucorticoid regulation of neuropeptide mRNAs in the rat striatum, 307
Chistakos, S., see Sonnenberg, J.L., 179
Chu, C., see Sonnenberg, J.L., 179
Chua, Jr., S.C., Leibel, R.L. and Hirsch, J., Food deprivation and age modulate neuropeptide gene expression in the murine hypothalamus and adrenal gland, 95
Clegg, K., see Nishimura, R.N., 39
Cole, R., see Nishimura, R.N., 39
Cortés, R., see Ceccatelli, S., 57
Coscia, C.J., Szücs, M., Barg, J., Belcheva, M.M., Bem, W.T., Khoobehi, K., Donnigan, T.A, Juszczak, R., McHale, R.J., Hanley, M.R. and Barnard, E.A., A monoclonal anti-idiotypic antibody to μ and δ opioid receptors, 299

Dahl, S.G., see Edvardsen, O., 31
Darlison, M.G., see Bateson, A.N., 333
Denêmes, D., Moniot, B., Lomri, N.-e., Thomasset, M. and Sans, A., Detection of calbindin-D 28k mRNA in rat vestibular ganglion neurons, 153
DiGiorganni, J., see Uhl, G.R., 23
Dickson, J.G., see Caviccholi, L., 319
Donnigan, T.A., see Coscia, C.J., 299
Douglass, J., Grimes, L., Shook, J., Lee, P.H.K. and Hong, J.-S., Systemic administration of kainic acid differentially regulates the levels of prodynorphin and proenkephalin mRNA and peptides in the rat hippocampus, 79
Dunnett, S.B., see Sirinathsinghji, D.J.S., 263
Dwyer, B.E., see Nishimura, R.N., 39

Edvardson, O. and Dahl, S.G., Molecular structure and dynamics of serotonin, 31
Emson, P.C., see Augood, S.J., 341
Emson, P.C., see Kiyama, H., 353
Emson, P.C., see Kiyama, H., 87

Faull, R.L.M., see Augood, S.J., 341
Finch, C.E., see Poirier, J., 191
Flanigan, T.P., see Caviccholi, L., 319
Forties, L.-P., Tremblay, J.P., Rafrafi, J. and Hawkes, R., A monoclonal antibody to conotoxin reveals the distribution of a subset of calcium channels in the rat cerebellar cortex, 209
Frantz, G.D., see Sonnenberg, J.L., 179
Freese, E., see Mill, J.F., 197
Fusco, M., see Caviccholi, L., 319

Gall, C., Murray, K. and Isackson, P.J., Kainic acid-induced seizures stimulate increased expression of nerve growth factor mRNA in rat hippocampus, 113
Glencorse, T.A., see Bateson, A.N., 333
Grima, E.A., see Sommerville, M.J., 1
Grimes, L., see Douglass, J., 79

Haleem-Smith, H., see Mill, J.F., 197
Hall, A.K. and Norman, A.W., Vitamin D-independent expression of chick brain calbindin-D_{28K}, 9
Hall, A.K., Developmental regulation of thymosin $\beta 10$ mRNA in the human brain, 175
Hanley, M.R., see Coscia, C.J., 299
Harbuz, H., Russel, J.A., Sumner, B.E.H., Kawata, M. and Lightman, S.L., Rapid changes in the content of proenkephalin A and corticotrophin releasing hormone mRNAs in the paraventricular nucleus during morphine withdrawal in urethane-anaesthetized rats, 285
Haroutunian, V., see Parks, K.M., 125
Harrison, P.J., Barton, A.J.L., Najlerahim, A., McDonald, B. and Pearson, R.C.A., Increased muscarinic receptor messenger RNA in Alzheimer's disease temporal cortex demonstrated by in situ hybridization histochemistry, 15
Harrison, P.J., Procter, A.W., Barton, A.J.L., Lowe, S.L., Najlerahim, A., Bertolucci, P.H.F., Bowen, D.M. and Pearson, R.C.A., Terminal coma affects messenger RNA detection in post mortem human temporal cortex, 161
Harvey, R.J., see Bateson, A.N., 333
Hawkes, R., see Forties, L.-P., 209
Heick, A., see Sonnenberg, J.L., 179
l'Héreault, S. and Barden, N., Monoaminergic regulation of proopiomelanocortin messenger RNA concentrations in primary cell cultures of rat hypothalamus, 327
Hess, M., see Poirier, J., 191
Hicks, A.A., see Bateson, A.N., 333
Higgins, G.A. and Olschowka, J.A., Induction of interleukin-1β mRNA in adult rat brain, 143
Hill, D.F., see Ojeda, S.R., 47
Hilt, D.C., see Lad, R.P., 239
Hirsch, J., see Chua Jr., S.C., 95
Hökfelt, T., see Ceccatelli, S., 57
Hong, J.-S., see Douglass, J., 79
Hunt, S.P., see Bateson, A.N., 333

Isackson, P.J., see Benson, D.L., 169
Isackson, P.J., see Gall, C., 113

Joh, T.H., see Kim, K.S., 277
Jones, E.G., see Benson, D.L., 169
Juszczak, R., see Coscia, C.J., 299

Katz, H., see Ojeda, S.R., 47
Kawata, M., see Harbuz, M., 285
Khoobehi, K., see Coscia, C.J., 299
Kim, K.S., Wessel, T.C., Stone, D.M., Carver, C.H., Joh, T.H. and Park, D.H., Molecular cloning and characterization of cDNA encoding tryptophan hydroxylase from rat central serotonergic neurons, 277
King, R., see Mill, J.F., 197
Kitayama, S., see Schaeffer, J.C., 271
Kiyama, H. and Emson, P.C., Colchicine induced expression of proneurotensin mRNA in rat striatum and hypothalamus, 353
Kiyama, H., McGowan, E.M. and Emson, P.C., Co-expression of cholecystokinin mRNA and tyrosine hydroxylase mRNA in populations of rat substantia nigra cells; a study using a combined radioactive and non-radioactive in situ hybridization procedure, 87
Kiyama, H., see Augood, S.J., 341
König, G., Salbaum, J.M., Wiestler, O., Lang, W., Schmitt, H.P., Masters, C.L. and Beyreuther, K., Alternative splicing

of the βA4 amyloid gene of Alzheimer's disease in cortex of control and Alzheimer's disease patients, 259
Kubo, T., see Saika, T., 157

Lad, R.P., Smith, M.A. and Hilt, D.C., Molecular cloning and regional distribution of rat brain cyclophilin, 239
Lahiri, D.K. and Robakis, N.K., The promotor activity of the gene encoding Alzheimer β-amyloid precursor protein (APP) is regulated by two blocks of upstream sequences, 253
Lamperti, E.D., Rosen, K.M. and Villa-Komaroff, L., Characterization of the gene and messages for vasoactive intestinal polypeptide (VIP) in rat and mouse, 217
Lang, W., see König, G., 259
Lee, P.H.K., see Douglass, J., 79
Lee, S., see Sonnenberg, J.L., 179
Leibel, R.L. see Chua Jr., S.C., 95
Leon, A., see Cavicchioli, L., 319
Li, X.-s. and Bartlett, W.P., Developmental expression of glial fibrillary acidic protein and glutamine synthetase mRNAs in normal and jimpy mice, 313
Lightman, S.L., see Harbuz, M., 285
Lin, C.-L., see Schaeffer, J.C., 271
Livett, B.G., see Wan, D.C.-C., 103
Livett, B.G., see Wan, D.C.-C., 135
Lomri, N.-e., see Denêmes, D., 153
Lowe, S.L., see Harrison, P.J., 161

Marley, P.D., see Wan, D.C.-C., 103
Marley, P.D., see Wan, D.C.-C., 135
Masters, C.L., see König, G., 259
Matsui, Y., see Sano, K., 149
Matsunaga, T., see Saika, T., 157
May, P.C., see Poirier, J., 191
McDonald, B., see Harrison, P.J., 15
McEwen, B.S., see Chao, H.M., 307
McGowan, E.M., see Kiyama, H., 87
McHale, R.J., see Coscia, C.J., 299
McLachlan, D.R.C., see Sommerville, M.J., 1
Mearow, K.M., see Mill, J.F., 197
Mill, J.F., Mearow, K.M., Purohit, H.J., Haleem-Smith, H., King, R. and Freese, E., Cloning and functional characterization of the rat glutamine synthetase gene, 197
Mohr, E. and Schmitz, E., Functional characterization of estrogen and glucocorticoid responsive elements in the rat oxytocin gene, 293
Moniot, B., see Denêmes, D., 153
Motoike, T., see Sano, K., 149
Murray, K., see Gall, C., 113

Najlerahim, A., see Harrison, P.J., 15
Najlerahim, A., see Harrison, P.J., 161
Nakamura, H., see Sano, K., 149
Nishimori, T., see Uhl, G.R., 23
Nishimura, R.N., Dwyer, B.E., Clegg, K., Cole, R. and Vellis, J. de, Comparison of the heat shock response in cultured cortical neurons and astrocytes, 39
Noguchi, K., see Saika, T., 157
Norman, A.W., see Hall, A.K., 9

O'Hara, B., see Uhl, G.R., 23
Ohtani, H., see Wada, C., 71
Ojeda, S.R., Hill, D.F. and Katz, H., The genes encoding nerve growth factor and its receptor are expressed in the developing female rat hypothalamus, 47
Okamoto, T. and Sumikawa, K., Antibiotics cause changes in the desensitization of ACh receptors expressed in *Xenopus* oocytes, 165
Olschowka, J.A., see Higgins, G.A., 143

Park, D.H., see Kim, K.S., 277
Parks, K.M., Sugar, J.E., Haroutunian, V., Bierer, L., Perl, D. and Wallace, W.C., Reduced in vitro phosphorylation of synapsin I (site 1) in Alzheimer's disease postmortem tissues, 125
Pearson, R.C.A., see Harrison, P.J., 15
Pearson, R.C.A., see Harrison, P.J., 161
Percy, M.E., see Sommerville, M.J., 1
Perl, D., see Parks, K.M., 125
Persson, H., see Srinivasan, M., 233
Piomelli, D., see Calignano, A., 347
Poirier, J., Hess, M., May, P.C. and Finch, C.E., Cloning of hippocampal poly(A) RNA sequences that increase after entorhinal cortex lesion in adult rat, 191
Procter, A.W., see Harrison, P.J., 161
Purohit, H.J., see Mill, J.F., 197

Rafrafi, J., see Forties, L.-P., 209
Robakis, N.K., see Lahiri, D.K., 253
Rosen, K.M., see Lamperti, E.D., 217
Russel, J.A., see Harbuz, M., 285

Sacktor, T.C., see Calignano, A., 347
Saika, T., Senba, E., Noguchi, K., Sato, M., Yoshida, S., Kubo, T., Matsunaga, T. and Tohyama, M., Effects of nerve crush and transection on mRNA levels for nerve growth factor receptor in the rat facial motoneurons, 157
Salbaum, J.M., see König, G., 259
Sano, K., Motoike, T., Nakamura, H., Matsui, Y., Teranishi, Y. and Takai, Y., Differential expression of the N-*myc*, c-*fos*, and *smg* p25A genes in human neuroblastoma cells during neuronal and Schwannian differentiation, 149
Sans, A., see Denêmes, D., 153
Sato, M., see Saika, T., 157
Schaeffer, J.C., Lin, C.-L., Kitayama, S. and Uhl, G.R., Ligand autoradiographic receptor screening. II. Expression of receptor cDNA in transfected COS cells grown on polyester disks and its recovery, 271
Scheartz, J.H., see Calignano, A., 347
Schmitt, H.P., see König, G., 259
Schmitz, E., see Mohr, E., 293
Selski, D.J., see Slemmon, J.R., 245
Senba, E., see Saika, T., 157
Shimada, S., see Uhl, G.R., 23
Shook, J., see Douglass, J., 79
Sirinathsinghji, D.J.S. and Dunnett, S.B., Increased proenkephalin mRNA levels in the rat neostriatum following lesion of the ipsilateral nigrostriatal dopamine pathway with 1-methyl-4-phenylpyridinium ion (MPP^+): reversal by embryonic nigral dopamine grafts, 263
Slemmon, J.R., Campbell, G.A., Selski, D.J. and Bramson, H.N., The amino terminus of the putative Drosophila choline acetyltransferase precursor is cleaved to yield the 67 kDa enzyme, 245
Smith, M.A., see Lad, R.P., 239
Sommerville, M.J., Percy, M.E., Bergeron, C., Yoong, L.K.K., Grima, E.A. and McLachlan, D.R.C., Localization and quantitation of 68 kDa neurofilament and superoxide dismutase-1 mRNA in Alzheimer brains, 1
Sonnenberg, J.L., Frantz, G.D., Lee, S., Heick, A., Chu, C., Tobin, A.J. and Chistakos, S., Calcium binding protein (calbindin-D_{28k}) and glutamate decarboxylase gene expression after kindling induced seizures, 179
Srinivasan, M., Yamamoto, Y., Brodin, E. and Persson, H., Chronic treatment with SCH-23390, a selective dopamine D_1 receptor blocker decreases preprotachykinin-A mRNA levels in nucleus tractus solitarii of the rabbit: role in respiratory control, 233
Stone, D.M., see Kim, K.S., 277
Sugar, J.E., see Parks, K.M., 125
Sumikawa, K., see Okamoto, T., 165
Sumner, B.E.H., see Harbuz, M., 285
Szücs, M., see Coscia, C.J., 299

Takai, Y., see Sano, K., 149
Teranishi, Y., see Sano, K., 149
Thomasset, M., see Denêmes, D., 153
Tobin, A.J., see Sonnenberg, J.L., 179
Tohyama, M., see Saika, T., 157
Toso, R.D., see Cavicchioli, L., 319
Tremblay, J.P., see Forties, L.-P., 209

Uhl, G.R., O'Hara, B., Shimada, S., Zaczek, R., DiGiorgianni, J. and Nishimori, T., Dopamine transporter: expression in *Xenopus* oocytes, 23
Uhl, G.R., see Schaeffer, J.C., 271

Vantini, G., see Cavicchioli, L., 319
Vellis, J. de, see Nishimura, R.N., 39
Villa-Komaroff, L., see Lamperti, E.D., 217

Wada, C. and Ohtani, H., Molecular cloning of rat cerebellin-like protein cDNA which encodes a novel membrane-associated glycoprotein, 71
Wallace, W.C., see Parks, K.M., 125
Walsh, F.S., see Cavicchioli, L., 319
Wan, D.C.-C., Marley, P.D. and Livett, B.G., Coordinate and differential regulation of proenkephalin A and PNMT mRNA expression in cultured bovine adrenal chromaffin cells: responses to secretory stimuli, 103
Wan, D.C.-C., Marley, P.D. and Livett, B.G., Coordinate and differential regulation of proenkephalin A and PNMT mRNA expression in cultured bovine adrenal chromaffin cells: responses to cAMP elevation and phorbol esters, 135
Wessel, T.C., see Kim, K.S., 277
Wiestler, O., see König, G., 259
Wisden, W., see Bateson, A.N., 333
Yamamoto, Y., see Srinivasan, M., 233
Yoong, L.K.K., see Sommerville, M.J., 1
Yoshida, S., see Saika,T., 157

Zaczek, R., see Uhl, G.R., 23